

Difference–restriction algebras with operators

Célia Borlido¹, Ganna Kudryavtseva², and Brett McLean^{3*}

¹ Universidade de Coimbra, CMUC, Departamento de Matemática, Coimbra, Portugal
`cborlido@mat.uc.pt`

² Faculty of Mathematics and Physics / Institute of Mathematics, Physics and Mechanics, University of Ljubljana

`ganna.kudryavtseva@fmf.uni-lj.si`

³ Department of Mathematics: Analysis, Logic and Discrete Mathematics, Ghent University
`brett.mclean@ugent.be`

Abstract

We exhibit an adjunction between a category of abstract algebras of partial functions that we call difference–restriction algebras and a category of Hausdorff étale spaces. Difference–restriction algebras are those algebras isomorphic to a collection of partial functions closed under relative complement and domain restriction; the morphisms are the homomorphisms. Our adjunction generalises the adjunction between the generalised Boolean algebras and the category of Hausdorff spaces. We define the finitary compatible completion of a difference–restriction algebra, and show that the monad induced by our adjunction yields the finitary compatible completion of any difference–restriction algebra. The adjunction restricts to a duality between the finitarily complete difference–restriction algebras and the locally compact zero-dimensional Hausdorff étale spaces, generalising the duality between Boolean algebras and locally compact zero-dimensional Hausdorff spaces. We then extend these adjunction, duality, and completion results to difference–restriction algebras equipped with arbitrary additional compatibility preserving operators.

Introduction

The study of algebras of partial functions is an active area of research that investigates collections of partial functions and their interrelationships from an algebraic perspective. The partial functions are treated as abstract elements that may be combined algebraically using various natural operations such as composition, domain restriction, ‘override’, or ‘update’. In pure mathematics, algebras of partial functions arise naturally as structures such as inverse semigroups, pseudogroups, and skew lattices. In theoretical computer science, they appear in the theories of finite state transducers, computable functions, deterministic propositional dynamic logics, and separation logic. Many different selections of operations have been considered, each

*Borlido was partially supported by the Centre for Mathematics of the University of Coimbra - UIDB/00324/2020, funded by the Portuguese Government through FCT/MCTES and partially supported by the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation program (grant agreement No. 670624). McLean was supported by the FWO Senior Postdoctoral Fellowship 1280024N.

leading to a different category of abstract algebras (see [8, §3.2] for a guide). Recently, dualities for some of these categories have started to appear [6, 5, 7, 9, 1], opening the way for these algebras to be studied via their duals.

In [2] and [3], we initiated a project to develop a general and modular framework for producing and understanding dualities for such categories. For this we are inspired strongly by Jónsson and Tarski’s theory of Boolean algebras with operators [4] and the duality between them and descriptive general frames. Our central thesis is that in our case the appropriate base class—the analogue of Boolean algebras—must be more than just a class of *ordered* structures but must record additional *compatibility* data. This reflects the fact that the union of two partial functions is not always a function.

In [2] and [3], we investigated algebras of partial functions for a signature we believe provides the necessary order and compatibility structure. The signature has two operations: set-theoretic *relative complement* and a *domain restriction* operation \triangleright given by: $f \triangleright g := \{(x, y) \in X \times Y \mid x \in \text{dom}(f) \text{ and } (x, y) \in g\}$. In [2], we gave and proved a finite equational axiomatisation for the class of isomorphs of such algebras of partial functions [2, Theorem 5.7]. We will refer to the algebras in this class as *difference–restriction algebras*. In [3], we gave a ‘discrete’ duality between the *atomic* difference–restriction algebras and a category of set quotients.

The main results of the present work are the elaboration of an adjunction between the category of difference–restriction algebras and the category of Hausdorff étale spaces (Thm. 2) and the extension of that theorem to algebras with additional operators (Thm. 6). We also show the monad induced by the adjunction gives a form of finitary completion of algebras (Thm. 3/Cor. 7(i)) and the adjunction restricts to a duality between the finitarily complete algebras and the locally complete zero-dimensional Hausdorff étale spaces (Thm. 4/Cor. 7(ii)).

Difference–restriction algebras and adjunction

An **algebra of partial functions** of the signature $\{-, \triangleright\}$ is a $\{-, \triangleright\}$ -algebra whose elements are partial functions from some (common) set X to some (common) set Y , and the interpretation of $-$ is *relative complement* and the interpretation of \triangleright is *domain restriction*.

A **difference–restriction algebra** is an algebra \mathfrak{A} of the signature $\{-, \triangleright\}$ that is isomorphic to an algebra of partial functions. We denote by **DRA** the category whose objects are difference–restriction algebras and whose morphisms are homomorphisms of $\{-, \triangleright\}$ -algebras. The operation $a \cdot b := (a - (a - b))$ gives difference–restriction algebras a semilattice structure.

An **étale space** is a surjective local homeomorphism $\pi: X \rightarrow X_0$, and π is **Hausdorff** if X is Hausdorff. A partial function $\varphi: X \rightarrow Y$ is **continuous** if when $V \subseteq Y$ is open in Y then $\varphi^{-1}(V)$ is open in X , and φ is **proper** if whenever $V \subseteq Y$ is compact then $\varphi^{-1}(V)$ is compact.

Definition 1. We denote by **HausEt** the category whose objects are Hausdorff étale spaces $\pi: X \rightarrow X_0$, and where a morphism from $\pi: X \rightarrow X_0$ to $\rho: Y \rightarrow Y_0$ is a continuous and proper partial function $\varphi: X \rightarrow Y$ satisfying the following conditions:

- (Q.1) φ **preserves equivalence**: if both $\varphi(x)$ and $\varphi(x')$ are defined, then $\pi(x) = \pi(x') \implies \rho(\varphi(x)) = \rho(\varphi(x'))$,
- (Q.2) φ is **fibrewise injective**: for every $(x_0, y_0) \in \tilde{\varphi}$, the restriction and co-restriction of φ induces an injective partial map $\varphi_{(x_0, y_0)}: \pi^{-1}(x_0) \rightarrow \rho^{-1}(y_0)$,
- (Q.3) φ is **fibrewise surjective**: for every $(x_0, y_0) \in \tilde{\varphi}$, the induced partial map $\varphi_{(x_0, y_0)}$ is surjective (that is, the image of $\varphi_{(x_0, y_0)}$ is the whole of $\rho^{-1}(y_0)$).

Theorem 2. *There exist adjoint functors $F: \mathbf{DRA} \rightarrow \mathbf{HausEt}^{\text{op}}$ and $G: \mathbf{HausEt}^{\text{op}} \rightarrow \mathbf{DRA}$.*

Roughly, F is ‘maximal filters’ and G is ‘partial sections with compact image’.

Duality and completion

Two elements of a difference–restriction algebra are **compatible** if $a_1 \triangleright a_2 = a_2 \triangleright a_1$ (corresponding to partial functions agreeing on their shared domain). The algebra is **finitarily compatibly complete** provided it has joins of each finite set of pairwise-compatible elements.

A **finitary compatible completion** of a difference–restriction algebra \mathfrak{A} is an embedding $\iota: \mathfrak{A} \hookrightarrow \mathfrak{C}$ of $\{-, \triangleright\}$ -algebras such that \mathfrak{C} is a difference–restriction algebra and finitarily compatibly complete and $\iota[\mathfrak{A}]$ is finite-join dense in \mathfrak{C} .

Theorem 3. *For each difference–restriction algebra \mathfrak{A} , the homomorphism $\eta_{\mathfrak{A}}: \mathfrak{A} \rightarrow (G \circ F)(\mathfrak{A})$ is the finitary compatible completion of \mathfrak{A} , where η is the unit of the adjunction of Theorem 2.*

We write $\mathbf{C}_{\text{fin}}\mathbf{DRA}$ for the full subcategory of \mathbf{DRA} consisting of the difference–restriction algebras that are finitarily compatibly complete. We write $\mathbf{Stone}^+\mathbf{Et}$ for the full subcategory of \mathbf{HausEt} consisting of the $\pi: X \rightarrow X_0$ such that X is locally compact and zero-dimensional.

Proposition 4. *The adjunction restricts to a duality between $\mathbf{C}_{\text{fin}}\mathbf{DRA}$ and $\mathbf{Stone}^+\mathbf{Et}$.*

Adjunction for difference–restriction algebras with operators

An n -ary operation Ω on \mathfrak{A} is **compatibility preserving** if whenever a_i, a'_i are compatible, for all i , we have that $\Omega(a_1, \dots, a_n)$ and $\Omega(a'_1, \dots, a'_n)$ are compatible, and Ω is an **operator** if it is *normal* ($\Omega(a_1, \dots, a_{i-1}, 0, a_{i+1}, \dots, a_n) = 0$) and is *additive* in each argument.

Let σ be a functional signature (disjoint from $\{-, \triangleright\}$).

The category $\mathbf{DRA}(\sigma)$ has *objects*: algebras of the signature $\{-, \triangleright\} \cup \sigma$ whose $\{-, \triangleright\}$ -reduct is a difference–restriction algebra, and such that the symbols of σ are interpreted as compatibility preserving operators, and *morphisms*: homomorphisms of $(\{-, \triangleright\} \cup \sigma)$ -algebras.

Let $\pi: X \rightarrow X_0$ be a Hausdorff étale space and R an $(n+1)$ -ary relation on X . The **compatibility relation** $C \subseteq X \times X$ is given by xCy if and only if $\pi(x) = \pi(y) \implies x = y$. Then R has the **compatibility property** if given $x_1Cx'_1, \dots, x_nCx'_n$ and $Rx_1 \dots x_{n+1}$ and $Rx'_1 \dots x'_{n+1}$, we have $x_{n+1}Cx'_{n+1}$.

Given subsets S_1, \dots, S_n of X , define $\Omega_R(S_1, \dots, S_n)$ by $\Omega_R(S_1, \dots, S_n) := \bigcup_{x_1 \in S_1, \dots, x_n \in S_n} \{x_{n+1} \in X \mid R x_1 \dots x_{n+1}\}$. The relation R **respects the topology** of $\pi: X \rightarrow X_0$ if whenever $S_1, \dots, S_n \subseteq X$ are compact open sets, then $\Omega_R(S_1, \dots, S_n)$ is a compact open set.

Take a partial function $\varphi: X \rightarrow Y$ and $(n+1)$ -ary relations R_X and R_Y on X and Y . Then φ satisfies the **reverse forth condition** if whenever $R_X x_1 \dots x_{n+1}$ and $\varphi(x_1), \dots, \varphi(x_n)$ are defined, then $\varphi(x_{n+1})$ is defined and $R_Y \varphi(x_1) \dots \varphi(x_{n+1})$. The partial map φ satisfies the **back condition** if whenever $\varphi(x_{n+1})$ is defined and $R_Y y_1 \dots y_n \varphi(x_{n+1})$, then there exist $x_1, \dots, x_n \in \text{dom}(\varphi)$ such that $\varphi(x_1) = y_1, \dots, \varphi(x_n) = y_n$ and $R_X x_1 \dots x_{n+1}$.

Definition 5. The category $\mathbf{HausEt}(\sigma)$ has *objects*: the objects of \mathbf{HausEt} equipped with, for each $\Omega \in \sigma$, an $(n+1)$ -ary relation R_Ω that has the compatibility property and respects the topology, where n is the arity of Ω , and *morphisms*: morphisms of \mathbf{HausEt} that satisfy the reverse forth condition and the back condition with respect to R_Ω , for every $\Omega \in \sigma$.

Theorem 6. *There is an adjunction $F': \mathbf{DRA}(\sigma) \dashv \mathbf{HausEt}(\sigma)^{\text{op}} : G'$ that extends the adjunction $F \dashv G$ of Theorem 2 in the sense that the appropriate reducts of $F'(\mathfrak{A})$ and $G'(\pi: X \rightarrow X_0)$ equal $F(\mathfrak{A})$ and $G(\pi: X \rightarrow X_0)$, respectively.*

Corollary 7. (i) *For every algebra \mathfrak{A} in $\mathbf{DRA}(\sigma)$, the embedding $\eta_{\mathfrak{A}}: \mathfrak{A} \hookrightarrow (G' \circ F')(\mathfrak{A})$ is the finitary compatible completion of \mathfrak{A} .*

(ii) *There is a duality between the categories $\mathbf{C}_{\text{fin}} \mathbf{DRA}(\sigma)$ and $\mathbf{Stone}^+ \mathbf{Et}(\sigma)^{\text{op}}$.*

References

- [1] Andrej Bauer, Karin Cvetko-Vah, Mai Gehrke, Samuel J. van Gool, and Ganna Kudryavtseva, *A non-commutative Priestley duality*, *Topology and its Applications* **160** (2013), no. 12, 1423–1438.
- [2] Célia Borlido and Brett McLean, *Difference–restriction algebras of partial functions: axiomatisations and representations*, *Algebra Universalis* **83** (2022), no. 3.
- [3] Célia Borlido and Brett McLean, *Difference–restriction algebras of partial functions with operators: discrete duality and completion*, *Journal of Algebra* **604** (2022), 760–789.
- [4] Bjarni Jónsson and Alfred Tarski, *Boolean algebras with operators. Part I*, *American Journal of Mathematics* **73** (1951), no. 4, pp. 891–939.
- [5] Ganna Kudryavtseva and Mark V. Lawson, *A perspective on non-commutative frame theory*, *Advances in Mathematics* **311** (2017), 378–468.
- [6] Mark V. Lawson, *A noncommutative generalization of Stone duality*, *Journal of the Australian Mathematical Society* **88** (2010), no. 3, 385–404.
- [7] Mark V. Lawson and Daniel H. Lenz, *Pseudogroups and their étale groupoids*, *Advances in Mathematics* **244** (2013), 117–170.
- [8] Brett McLean, *Algebras of partial functions*, Ph.D. thesis, University College London, 2018.
- [9] Brett McLean, *A categorical duality for algebras of partial functions*, *Journal of Pure and Applied Algebra* **225** (2021), no. 11, 106755.